Reconstructed neurtino energy

Danaisis Vargas

Institute of High Energy Physics - IFAE

dvargas@ifae.es

October 26, 2017

Introduction
T2K and NEUT

Using the NEUT simulation code, the reaction studied was:

- $\nu_\mu \rightarrow {}^{12}C$

CCQE or CC0π



T2K



Figure 4.8: Schematic representation of the T2K configuration.

- $L_{near} = 280$ m;
- $L_{far} = 295$ km.

Theoretical Model

Equations for CCQE

Reconstructed neutrino energy (E_ν^{rec}):

$$E_\nu^{rec} = \frac{2(M_n - E_B)E_\mu - (E_B^2 + M_\mu^2 - 2M_nE_B + \Delta M^2)}{2(M_n - E_B - E_\mu + |\vec{k_\mu}|cos\theta_\mu)} \quad (1)$$

$$\Delta M^2 = M_n^2 - M_p^2$$

$$E_\mu = \sqrt{|\vec{k_\mu}|^2 + M_\mu^2}$$

Transfered momentum (Q^2_{rec}):

$$Q^2_{rec} = 2\ E_\nu^{rec}\ (E_\mu - |\vec{k_\mu}|cos\theta_\mu) - M_\mu^2 \quad (2)$$

- $M_n = 939.565379$ MeV;
- $M_p = 938.272046$ MeV;
- $M_\mu = 105.6583715$ MeV;
- $E_B = 24$ MeV.

Theoretical Model

Equations for oscillation probability

Probability of disappearance of the ν_μ:

$$P_{(\nu_\mu \to \nu_\mu)} = 1 - sen^2\theta_{23}\ sen^2\left(1.267\ \frac{\Delta M^2_{32}\ L_{far}}{E^{rec}_\nu}\right) \quad (3)$$

- $sen^2(2\theta_{13}) = 0.14$;
- $sen^2\theta_{23} = 0.514$;
- $\Delta M^2_{32} = 2.51 \times 10^{-3}$ eV^2.

Important!!!



Results
Reconstructed neutrino energy



Figure 1: Reconstructed neutrino energy vs. muon angle cosine.

Results
Reconstructed neutrino energy



Figure 2: Reconstructed neutrino energy vs. transferred moment.

Results
Reconstructed neutrino energy



Figure 3: Reconstructed neutrino energy vs. transferred moment matrix with bins of 0.5 GeV for E_ν^{rec} and 0.5 GeV^2 for Q^2_{rec}.

Results
Reconstructed neutrino energy



Figure 4: No. of ν_{mu} events vs. $E_\nu^{true} - E_\nu^{rec}$ for each block of the matrix.

Results
Reconstructed neutrino energy



Figure 5: Reconstructed neutrino energy vs. muon angle.

Results
Reconstructed neutrino energy



Figure 6: Reconstructed neutrino energy vs. muon angle matrix with bins of 0.5 GeV for E_ν^{rec} and 20○ for θ_μ.

Results
Reconstructed neutrino energy



Figure 7: No. of ν_{mu} events vs. $E_\nu^{true} - E_\nu^{rec}$ for each block of the matrix.

Results
Reconstructed neutrino energy



Figure 8: No. of ν_{mu} events vs. $E_\nu^{true} - E_\nu^{rec}$ for each block of the matrix.

Results
Reconstructed neutrino energy



Figure 9: .

Results
Reconstructed neutrino energy



Figure 10: Bla bla bla.

Results
Reconstructed neutrino energy



Figure 11: Bla bla bla.

Results

ν_μ disappearance probability



Figure 12: ν_μ disappearance probability for different μ angles.

Thank you!!!